

**OKLAHOMA DEPARTMENT OF CORRECTIONS  
Periodic Physical Examination and Laboratory Requirements  
Male**

<b>Periodic Physical Examination and Laboratory Requirement: Male (5-ACI-6A-27)</b>		
<b>Age</b>	<b>Requirements</b>	<b>Frequency</b>
Periodic PE Ages 18-39 yrs.	<ul style="list-style-type: none"> <li>Physical Examination</li> </ul>	<ul style="list-style-type: none"> <li>Every 3 years</li> </ul>
Periodic PE Ages 40-64 yrs.	<ul style="list-style-type: none"> <li>Physical Examination</li> <li>Fasting Lipid Profile</li> <li>CMP</li> <li>Hepatitis C Profile (CPL 162)</li> <li>Fecal occult blood testing X 3 beginning age 50</li> <li>Baseline EKG - age 50</li> </ul>	<ul style="list-style-type: none"> <li>Every 1-3 years</li> <li>Every 1-3 years</li> <li>Every 1-3 years</li> <li>Baseline Only</li> <li>Annually Baseline, and then by provider discretion</li> </ul>
Periodic PE Ages 65 and older	<ul style="list-style-type: none"> <li>Physical Examination</li> <li>Fasting Lipid Profile</li> <li>CMP</li> <li>Fecal occult blood</li> </ul>	Annually Every 1-3 years to age 75, then provider discretion afterwards <ul style="list-style-type: none"> <li>Every 3 years</li> <li>Annually until age 75, and then by provider discretion</li> </ul>

**Physical Examination and Laboratory Requirements for Inmates Returning From GPS  
Male**

<b>GPS Physical Examination and Laboratory Requirement: Male</b>	
Initial PE Age 39 and younger	<ul style="list-style-type: none"> <li>Tuberculin skin test</li> <li>HIV testing – Enzyme-Linked Immunosorbent Assay (ELISA). The Western Blot method will be utilized if the baseline ELISA test result is reactive (or positive).</li> <li>Rapid Plasma Reagin (Syphilis)</li> <li>CMP</li> </ul>
Initial PE Ages 40-49 yrs.	<ul style="list-style-type: none"> <li>CMP</li> <li>Tuberculin skin test</li> <li>HIV testing – Enzyme-Linked Immunosorbent Assay (ELISA). The Western Blot method will be utilized if the baseline ELISA test result is reactive (or positive).</li> <li>Rapid Plasma Reagin (Syphilis)</li> </ul>
Initial PE Ages 50-64 yrs.	<ul style="list-style-type: none"> <li>CMP</li> <li>Tuberculin skin test</li> <li>HIV testing – Enzyme-Linked Immunosorbent Assay (ELISA). The Western Blot method will be utilized if the baseline ELISA test result is reactive (or positive).</li> <li>Rapid Plasma Reagin (Syphilis)</li> </ul>
Initial PE Ages 65 and older	<ul style="list-style-type: none"> <li>CMP</li> <li>Tuberculin skin test</li> <li>HIV testing – Enzyme-Linked Immunosorbent Assay (ELISA). The Western Blot method will be utilized if the baseline ELISA test result is reactive (or positive).</li> <li>Rapid Plasma Reagin (Syphilis)</li> </ul>