MSRM 140117.01.12.4 (R-4/19)

OKLAHOMA DEPARTMENT OF CORRECTIONS NURSING PRACTICE PROTOCOLS

Percutaneous Endoscopic Gastrostomy Tube (PEG)

<u>ALL</u> PROCEDURES MUST HAVE A CORRESPONDING ORDER DOCUMENTED IN THE INMATES EHR. IF THERE IS NO DOCUMENTED ORDER, WRITE A VERBAL ORDER AND ASSIGN TO THE PROVIDER FOR SIGNATURE. CMA'S THAT HOLD A CURRENT GASTRO CERTIFICATION MAY USE THIS NURSING PROTOCOL.

Subjective Data:		Allergies:			
Chief complaint:					
Peg Materials:					
Catheter Tip Syringe A(60cc) Stethoscope Gloves Water Medical Tape		Cotton-tip swab (to clean under anchor) Ointment (if ordered) Gauze dressing (if drainage around tube) Formula			
Objective Data: (clinically indicated VS)					
BP	Pulse	Resp	Temp		Vt
Urine:	□ Clear	☐ Dark [Cloudy	□ Bloody	☐ Foul oder
CMA's ARE TO CONTACT THE RN IMMEDIATELY. THE RN IS TO CONTACT HEALTH CARE PROVIDER: Health care provider must be called if not on site or if after clinic hours. □ Bloody or coffee colored drainage through tube □ Stomach becomes bloated or swollen and tight □ Temperature of 100.4 degrees F. □ Peg tube comes out or is shorter than when it was put in. □ Coughing or vomiting during/after feeding □ S/S if infection at insertion site (redness, warm to touch, firm to touch, tender) Health Care Provider: □ Time Notified: □ Orders Received for Treatment: □ Yes □ No If physical exam is negative for any of the above s/s and/or there is no need for additional medication/treatment, proceed with nursing					
Plan: Interventions: (check all that apply)					
□ Peg Tube Feeding □ Routine Peg Care					
 □ Provided privacy and procedure explained to inmate. □ Hands washed with soap and water and gloves applied. □ Inmate placed in semi-fowlers position. □ Peg adapter cleanse with alcohol pad. □ Peg tube placement assessed with auscultation. Injected 30 cc to 50 cc of air with syringe. Whooshing/gurgling sound present. □ Aspirated stomach contents for residual. Residual amount obtained cc. Instilled aspirated contents back into feeding tube. (Notify health care provider if gastric residual > 50 cc.) □ Peg tube flushed with 30 cc water to clear tube of gastric fluid. □ Formula (name and amount) administered. □ Peg tube flushed with 30 cc water after feeding. □ C free water given. □ No signs/symptoms of bloating, nausea, or vomiting during/after feeding. □ Gloves removed, disposed, and hands rewashed. □ Inmate tolerated procedure well. □ Education/Intervention: Instructed patient to follow-up sick call if experiencing any signs and symptoms that warrant treatment. 					and gloves applied. d with soap and water. e. ape. d. ds rewashed.
Inmate verbalizes understanding of instructions.					
Progress Note: Health Care Provider Signature/Credentials: RN/LPN/CMA Signature/Credentials:				Date:	Time:
Inmate Name (Last, First)					