

OTTAWA COUNTY

DRUG OVERDOSE DEATHS¹ (2017-2021)

NUMBER OF UNINTENTIONAL DRUG OVERDOSE DEATHS BY TYPE OF SUBSTANCE*

28

total number of unintentional drug overdose deaths

20

Methamphetamine

8

Other Opioids (excluding fentanyl)

5

Fentanyl

0

Cocaine

0

Alcohol

<5

Benzodiazepines

* Categories are not mutually exclusive. Deaths may involve more than one substance.

From 2017 to 2021, **Ottawa County** had the **36th highest** drug overdose death rate in the state.

Adults aged 45-54 years had the highest rate of death.




61% of deaths occurred at a home or apartment.

NON-HISPANIC WHITE Ottawa County residents had the **highest** rate of death.

- Males were **3.8 times more likely** to die than females.
- Drug overdose **death rates decreased 1%** from 18.1 per 100,000 (2012-2016) to 18 per 100,000 (2017-2021).
- **One in four (25%)** people who died had a history of mental health problems.
- **More than two out of three (68%)** people who died had a history of substance use.

UNINTENTIONAL DRUG OVERDOSE DEATH RATES¹ BY COUNTY OF RESIDENCE²

► OKLAHOMA, 2017-2021



¹ Some rates are based on small numbers and may be unstable.

² County of residence was unknown for 42 decedents.

Data Source: Fatal Unintentional Poisoning Surveillance System, OSDH Injury Prevention Service

OTTAWA COUNTY

DRUG OVERDOSE HOSPITALIZATIONS[†] (2019-2021)

Ottawa County had the **34th** highest nonfatal drug overdose hospitalization rate in the state.



ADULTS AGED **35-44** had the highest hospitalization rate.



There were **80** Ottawa County Residents hospitalized for a nonfatal drug overdose from 2019-2021.

- **45%** of hospitalizations were unintentional and **54%** due to self harm.
- Females were **31% more likely** to be hospitalized than males.
- Drug overdose **hospitalization rates decreased 24%** from 113.4 per 100,000 (2016-2018) to 86.6 per 100,000 (2019-2021).
- Non-Hispanic White Ottawa County residents had the **highest rate** of hospitalizations.


The most common substances involved in hospitalizations were:

- ▶ Opioids
- ▶ Antidepressants
- ▶ Non-opioid pain medications
- ▶ Stimulants
- ▶ Antipsychotics

[†] Includes all intents of nonfatal drug overdose – unintentional, self harm, undetermined, and assault

NONFATAL DRUG OVERDOSE HOSPITALIZATION RATES³ BY COUNTY OF RESIDENCE⁴

▶ OKLAHOMA, 2019-2021⁵



³ Some rates are based on small numbers and may be unstable.

⁴ County of residence was unknown for 64 patients.

⁵ Preliminary 2021 data

Data Source: OSDH Center for Health Statistics, Oklahoma Hospital Discharge Data
Compiled by: OSDH Injury Prevention Service

