



OKLAHOMA STATE
DEPARTMENT OF HEALTH

State of the County's Health Report

Working Together For Health

Ellis County

Oklahoma ranks near the bottom in many key health status indicators. Most of these outcomes are related to conditions that Oklahomans must live with every day. Poverty, lack of insurance, limited access to primary care, and inadequate prenatal care, along with associated risky health behaviors (low fruit/vegetable consumption, low physical activity, a high prevalence of smoking) all contribute to the poor health status of our citizens.

The vision of the Oklahoma Health Improvement Plan (OHIP) involves local partnerships and communities working together to improve and sustain the physical, social, and mental well-being of all people in Oklahoma. The targeted flagship initiatives of children's health improvement, tobacco use prevention, and obesity reduction will help to maximize opportunities for all Oklahomans to lead healthy lives. If you would like more information about OHIP, please visit the Oklahoma State Department of Health website at http://www.ok.gov/health/Organization/Board_of_Health/OHIP.html.

This report focuses on health factors and demographics in Ellis County. Awareness and thoughtful application of this health data can assist us in our joint endeavors to improve the health status of our local citizens.



Together We Can Move Forward

Inside This Issue

Table of Contents	1	Tobacco Use Prevention	7
County Demographics	2	Healthy People 2010 Table	8
Top 10 Leading Causes of Death	2	Health Care Costs Summary	9
Top 10 Leading Causes of Death Table	3	County Health Department Usage	10
Nutrition & Overweight	4	Health Education	11
Physical Activity & Fitness	4	Primary Care Map	11
Diabetes	5	Board of Health Map	12
Teen Births	5	Oklahoma Health Improvement Plan	12
Infant Mortality	6	Reference List	13
Low Birth Weight	6	Turning Point	14
Injury & Violence	7	Contact Information	14

County Demographics

Population estimates¹

- 9.5% decrease from 1990 to 2000 (4,469 to 4,043)
- 1.8% decrease from 2000 to 2008 (4,043 to 3,971)
- Ranked 58th for growth in state

2008 Census Estimates²


- Hispanic/Latino ethnicity = 4.1%
- Race
 - Whites = 98%
 - Native Americans = 1.5%
 - Blacks = 0.5%
- Age
 - Under 5 = 6.9%
 - 65 and over = 21.3%
 - Median age = 45.3 years

2000 Census³

- Housing units
 - Occupied = 1,769 (82%)
 - Vacant = 377 (18%)

- Disability (ages 21 to 64) = 23.7%
 - state = 21.5% national = 19.2%
 - 61.3% of disabled (ages 21-64) are employed
- Individuals below poverty = 12.5%
 - state = 14.7% national = 12.4%
- Families below poverty = 9.2%
 - state = 11.2% national = 9.2%

1990, 2000, & 2008 Population by Age Groups, Ellis County




Top 10 Leading Causes of Death


The top 10 leading causes of death⁴ table on the next page displays a broad picture of the causes of death in Ellis County. Since many health-related issues are unique to specific ages, this table provides causes of death by age group at a glance. The causes of death that are present across almost every age group have been highlighted. From 1983 to 1993 heart disease killed 1,129 people in Ellis County and is still the leading cause of death with all age groups combined.

From 2006-2008, the total charges for all heart disease-related hospitalizations for Ellis County residents was \$8,168,247.00, which translated to an average of \$30,592.69 per discharge.⁵ Total charges are an indicator of community health, however, these are hospital-based charges and not payments or costs of actual illness. The direct and indirect costs of disease are actually much higher (i.e., loss of employee wages, loss of tax revenue, loss of years of life).

Age-Adjusted Death Rates by County, OSDH Vital Statistics, 2002-2006⁴



Heart Disease Death Rates by Demographic Groups, Ellis County, 2002-2006⁴



* Age-Adjusted Death Rate per 100,000 Population

Legend

- 759.5 - 885.4
- 885.5 - 970.8
- 970.9 - 1034.6
- 1034.7 - 1149.0
- Missing

State = 945.6
National = 810.1
* Note: Data classified by Quartiles

Top 10 Causes of Death by Age Group Ellis County 2002-2006

Rank	0-4	05-14	15-24	25-34	35-44	45-54	55-64	65+	All Ages
1	CANCER < 4	UNINTENT. INJURY < 4		UNINTENT. INJURY < 4	HEART DISEASE < 4	CANCER 4	CANCER 8	HEART DISEASE 80	HEART DISEASE 91
2				HEART DISEASE < 4	CANCER < 4	DIABETES MELLITUS < 4	HEART DISEASE 7	CANCER 29	CANCER 43
3					DIABETES MELLITUS < 4	UNINTENT. INJURY < 4	BRONCHITIS/ EMPHYSEMA/ ASTHMA < 4	STROKE 19	STROKE 21
4					STROKE < 4	HEART DISEASE < 4	MENINGITIS < 4	BRONCHITIS/ EMPHYSEMA/ ASTHMA 18	BRONCHITIS/ EMPHYSEMA/ ASTHMA 20
5					UNINTENT. INJURY < 4	NEPHRITIS < 4	UNINTENT. INJURY < 4	DIABETES MELLITUS 11	DIABETES MELLITUS 15
6							DIABETES MELLITUS < 4	ALZHEIMER'S DISEASE 10	UNINTENT. INJURY 14
7							INFLUENZA/ PNEUMONIA < 4	INFLUENZA/ PNEUMONIA 9	ALZHEIMER'S DISEASE 10
8							STROKE < 4	NEPHRITIS 6	INFLUENZA/ PNEUMONIA 10
9							SUICIDE < 4	UNINTENT. INJURY 4	NEPHRITIS 7
10							VIRAL HEPATITIS < 4	TWO CAUSES TIED < 4	TWO CAUSES TIED < 4

Note: The numbers less than 4 have been shown as "< 4" to protect the privacy of the subjects
 Data source: Vital Statistics, Health Care Information Division, Oklahoma State Department of Health
 Produced by: Community Development Service, Oklahoma State Department of Health


Nutrition & Overweight

With obesity at epidemic levels, steps need to be taken to control this issue. Surveillance systems have improved and been expanded to obtain more accurate county-level data. While efforts continue on this front, communities must utilize this information to improve the problem itself. With health care costs being, on average, \$395 more for a person under the age of 65 who is obese than a person of the same age who is not obese⁶, estimated health care costs related to obesity for Ellis County soar to almost \$3.2 thousand. These costs only go up when the 65 and over population are included.

The Youth Risk Behavior Surveillance System (YRBS)⁷, while not producing county-level data, shows that in 2005, 2007, and 2009 combined, 15.8% of high school children participating in the survey classified themselves as overweight. Further, 14.7% classified themselves as obese. Interestingly, 8.9% of the males said they were obese compared to 5.8% of females. This trend was reversed in the overweight category with 8.0% of females and 7.8% of males, statewide. Although, neither weight category is statistically significant by gender.


According to the Behavioral Risk Factor Surveillance System (BRFSS, 2003 & 2005)⁸, 98.4% of Ellis County adults did not eat the recommended 5 servings of fruits and vegetables a day. Increasing fruit and vegetable consumption is an economical way to control many health-related issues such as obesity and diabetes.

Percent of Adults who are Overweight, BRFSS, 2003-2008⁸



* Note: Data classified by Quartiles


Percent of Adults who are Obese, BRFSS, 2003-2008⁸



* Note: Data classified by Quartiles

Physical Activity & Fitness

Percent of Adults with No Physical Activity Within Past Month, BRFSS, 2003-2008⁸



* Note: Data classified by Quartiles

The increasing inactivity of the U.S. population is contributing to an increase in numerous poor health-related outcomes. Physical inactivity robs the body of precious energy needed to function properly, in turn health declines, and rates of various chronic diseases escalate.


According to the 2003-2008 BRFSS⁸, it is estimated that 35.6% (1,090) of people in Ellis County had no leisure activity in the past month (at the time they were surveyed).

Statewide, 55% of high school students did not participate in physical activity for at least 60 minutes per day on five or more days in a week.⁷ Efforts need to be made to increase physical activity for our youth to build long-term healthy habits.

Diabetes

As stated previously, poor nutrition and lack of physical activity are linked to many chronic issues including diabetes. It is often hard to distinguish between Type I and Type II diabetes in large datasets but looking at the overall impact of diabetes is necessary to examine a community's health.

Percent of Adults Who Have Ever Been Told They Have Diabetes, BRFSS, 2004-2008⁸



* Note: Data classified by Quartiles

From 2006 to 2008, there were 0 hospital discharges for diabetes among Ellis County residents.

According to the 2004-2008 BRFSS, it is estimated that 6.7% (205) of Ellis County citizens have been diagnosed by a health professional as having diabetes.

In 2007, the per capita annual healthcare costs for people with diabetes was \$11,744 compared to \$5,106 for people without diabetes.⁹ Persons with diabetes accumulate an estimated \$2,717,741.70 in health care costs in one year for Ellis County. Actual hospital charges account for only 0.0% of the total health care impact of diabetes.

Teen Births


While births to teen mothers (age 15 to 19) have been on the decline in both the U.S. and Oklahoma, Oklahoma has moved down in the rankings according to the United Health Foundation.¹⁰ In 1993, the rate for teen births in the U.S. was 60.3 per 1,000 15-19 year old females and 67.1 in Oklahoma, ranking 35th in the country.¹⁰ In 2003, the rate decreased to 47.7 in the U.S. and 59.7 in Oklahoma, accounting for a decrease of 21% and 11%, respectively.¹⁰ However, while the rate decreased, Oklahoma continued to fall in the rankings (41st). In 2009, the rate of teen births stayed the same for Oklahoma (59.6) but the state ranked 45th in the country.¹⁰

Children of teen mothers are more likely to display poor health and social outcomes than those of older mothers, such as premature birth, low birth weight, higher rates of abuse and neglect, and are more likely to go into foster care or do poorly in school.¹¹

According to Oklahoma Vital Statistics, Ellis County had a teen birth rate of 70.7 in 2007, which accounted for a 17% increase from 2003 (60.6) and a 53% increase from 1993 (46.2)¹². The map represents a five-year average of teen birth rates, 2003-2007.¹²

On average in Oklahoma, births to teen mothers accumulate \$3,807 a year for each teenage birth¹³, which is often passed on to citizens. With an average of 6.0 births per year (2003-2007)¹², teen pregnancy costs the citizens of Ellis County \$19,200.00 a year.

Rate of Births to Teen Mothers (Age 15-19), OSDH Vital Statistics, 2003-2007¹²




* Note: Data classified by Quartiles

Infant Mortality

Protecting the most vulnerable populations is a task for every Oklahoman, socio-demographic variables have repeatedly been shown to affect infant mortality and birth weight. Income, education, race/ethnicity, access to medical care and social services, and social support are just a few of the characteristics that alter these rates, both positively and negatively.¹⁶ Differences in infant mortality rates are most apparent within racial categories. From 2003-2005 within the U.S., the infant mortality rate among white mothers was 5.7 deaths per 1,000 live births, while black or African American mothers had an infant mortality rate of 13.3 and American Indian mothers a rate of 8.4. Interestingly, Hispanic mothers had rates similar to white mothers.¹⁵

Infant Mortality Rate, Oklahoma Vital Statistics, 2002-2006⁴



State = 8.0 per 1,000 births
 National = 6.8 per 1,000 births (2003-2005)
 * Note: Data classified by Quartiles


Within Oklahoma, this trend continues. From 2002-2006, the infant mortality rate was highest for black or African American mothers (16.0), followed by American Indian (8.4), White (7.1), and Asian (5).⁴ The Hispanic infant mortality rate was 7.9.⁴

With an average of 412 infant deaths a year across Oklahoma⁴, it is difficult to obtain stable county-level data by race. From 2002-2006, the overall infant mortality rate for Ellis County was missing.

Low Birth Weight

Birth weight has been a long standing indicator of long-term health outcomes as birth weight has been linked to certain adult chronic conditions such as high blood pressure, Type II diabetes, and heart disease.¹⁷ Low birth weight is defined as any baby born weighing less than 2,500 grams or 5 pounds and 8 ounces (including very low birth weight = less than 1,500 grams/3 pounds, 5 ounces).¹⁷ Low birth weight is associated with premature births, multiple births, birth defects, chronic health problems or infections in the mother, smoking, alcohol or drug use, placental problems, inadequate maternal weight gain or socioeconomic factors.¹⁷ Understanding the trends behind low birth weight can assist in pinpointing causes specific to a community.

Percent of Live Births with Low and Very Low Birth Weight, OSDH Vital Statistics, 2003-2007¹²



In Oklahoma from 2003-2007, 8.1% of the live births were less than 2,500 grams. As with infant mortality, babies born to black or African American mothers have the highest rates of low birth weight (14.1%), followed by White (7.5%), American Indian and Asian (6.9% each).¹² This trend is similar to the national data.¹⁸

In Ellis County, from 2003-2007 the low birth weight rate was 6.7%.¹² This was a 17.3% decrease from the state and national low birth weight rate. Because of the small numbers that occur in this category, county level data by race is unavailable.

State = 8.1% of live births
 National = 8.1% of live births (2003-2006)
 Note: % of births with birth weight under 2500 grams
 Data classified by Quartiles

Injury and Violence


Across the nation and the state of Oklahoma, unintentional and violence-related injuries are on the rise. Unintentional injuries account for the 5th leading cause of death in the United States and Oklahoma for 2002-2006. For persons ages 1 to 44 in Oklahoma, unintentional injuries are the leading cause of death.⁴

This trend changes slightly in Ellis County. Unintentional injuries are in the top ten leading causes of death for ages 15 and over.

It is estimated that for every motor vehicle-related death \$1.3 million in economic costs are incurred (2008 data).¹⁶ For Ellis County, which has an average of 1.2 motor vehicle-related deaths a year⁴, the estimated economic costs are almost \$1.6 million a year.

Violence-related injuries (homicide and suicide) in Ellis County are ranked in the top 10 causes of death for persons from 55 to age 64 and suicide is the 9th leading cause of death for ages 55 to 64.⁴

Age-Adjusted Suicide Rate by County, OSDH Vital Statistics, 2002-2006⁴




Tobacco Use Prevention

According to the 2005 State of the State's Health Report¹⁹, tobacco use among Oklahomans has remained fairly stable from 1990 to 2002. The good news is that total cigarette sales in Oklahoma (tribal and non-tribal combined) have dropped from 98.2 packs per capita in fiscal year 2005 to 86.7 packs per capita during fiscal year 2008. The national average dropped during this same time period.²⁰

Tobacco use is no longer just the problem of the individual but also the community as a whole. With health care costs on the rise, targeting areas such as tobacco use is an effective way to control those costs.


Per Capita Cigarette Sales: Oklahoma & United States²⁰



the rise, targeting areas such as tobacco use is an effective way to control those costs. The Oklahoma Tobacco Helpline (1-800-QUIT-NOW), supported jointly by the Oklahoma Tobacco Settlement Endowment Trust and the Oklahoma State Department of Health, continues to experience high call volume. Over 37,800 Oklahomans received free cessation assistance through the Helpline in fiscal year 2009.²⁰ Since inception of the Helpline in August 2003, over 110,000 Oklahomans have received free cessation assistance.²⁰

The CDC estimated that a person who used tobacco accrued over \$3,300 in health care costs per year.²¹ According to the BRFSS (2005-2009)⁸, it is estimated that 18.9% (579) of adults in Ellis County use tobacco of some sort. Medical costs accumulated by those persons are over \$1.9 million a year for Ellis County.

Percent of Adults by Smoking Status, Ellis County, 2001-2009, Five-Year Averages⁸



Healthy People 2010 Table

Healthy People 2010 Indicators	Most Recent Data: Year(s)						2010 Target
	Ellis County		Oklahoma		United States		
Prevalence of Obese (Aged 18+)	2002-2008	22.3%	2008	31.0%	2008	26.7%	15%
No Leisure-Time Physical Activity (Aged 18+)	2002-2008	33.6%	2008	31.5%	2008	24.6%	20%
Prevalence of Smoking (Aged 18+)	2002-2008	27.9%	2008	24.7%	2008	18.4%	12%
Infant Mortality (Per 1,000 of births)	2002-2006	0	2006	8.1	2006	6.8	4.5
Low Birth Weight Infants (Percent of live births)	2002-2007	0.0%	2006	8.3%	2006	8.3%	5%
Very Low Birth Weight Infants (Percent of live births)	2002-2007	0.0%	2006	1.6%	2006	1.5%	0.9%
First Trimester Prenatal Care (Percent of births)	2002-2007	76.6%	2006	75.6%	2006	83.2%	90%
Prevalence of Diabetes (Aged 18+)	2002-2008	6.6%	2008	11.3%	2008	9.2%	2.5%
Lack of Health Insurance (Aged 18-64)	2002-2008	27.1%	2008	22.8%	2008	17.1%	0%
Prevalence of Binge Drinking (Aged 18+)	2002-2008	19.5%	2008	12.2%	2008	15.6%	6%
Coronary Heart Disease Death *	2002-2006	177.4	2006	184.5	2006	144.4	166.0
Cancer Death *	2002-2006	137.1	2006	194.9	2006	180.8	159.9
Unintentional Injury Death *	2002-2006	70.3	2006	55.6	2006	39.3	17.5
Transportation-Related Death *	2002-2006	38.7	2006	21.0	2006	14.5	9.2

Note: * means (Age-adjusted death per 100,000 to the 2000 U.S. standard population).

Reference:

- [1] Healthy People 2010 volume I and II, U.S. Department of Health and Human Services, November 2000.
- [2] Centers for Disease Control and Prevention (CDC), Wide-Ranging Online Data for Epidemiologic Research: Data for Oklahoma and United States.
- [3] CDC, National Center for Chronic Disease Prevention and Health Promotion, the Behavioral Risk Factor Surveillance System (BRFSS): Data for Oklahoma and United States.
- [4] Oklahoma State Department of Health (OSDH), OK2SHARE, BRFSS: Data for Oklahoma Counties.
- [5] OSDH, OK2SHARE, Vital Statistics: Data for Oklahoma Counties.

Health Care Cost Summary

Cardiovascular Disease (Heart Disease)

- Average hospital discharges per year⁵ = 89
- Average charges⁵ = \$30,592.69
- Total—\$2,722,749.00 a year

Obesity

- 26.4% of population⁸ (809)
- \$395 in additional medical costs per person aged 18-64⁶
- Total—\$319,555.00

Diabetes

- Data Not Available

Teen Pregnancy

- Average 6 births to females aged 15-19 a year¹²
- \$3,200 in costs a year¹³
- Total—\$19,200.00 a year

Motor Vehicle-Related Injury Death

- Average 1.2 deaths per year⁴
- \$1,300,000.00 in economic costs per death¹⁶
- Total—\$1,560,000.00 a year

Tobacco Use

- 18.9% of population⁸ (579)
- \$3,300 in health care costs²¹
- Total—\$1,910,700.00 a year


Grand Total for
Ellis County:

\$6,532,204.00



County Health Department Usage

County Health Department Unduplicated Clients, and Visits by Program, Ellis County, State Fiscal Year 2009



Data Note: Data is reflective of all services offered in a county, including county health departments and contracts.

■ Unduplicated Clients ■ Visits

ELLIS


Population-Based Services by Event Type, Ellis County, SFY09

Event Type	Number of Events	Total Attendees
Presentation/Class	12	809
Grand Total	12	809

Population-Based Services by Main Topic, Ellis County, SFY09

Topic	Number of Events	Total Attendees
General Health Department Services	2	85
Oral Health	8	679
Tobacco Education	2	45
Grand Total	12	809

Health Education



CATCH Kids Club Sites

- CKC Pilot Sites 2009-2010 (17)
- ★ NEW CKC Sites 2009-2010 (40)


**Ellis County
Health Educator**
 Jolena Graves
 7th & Oklahoma Ste. 9
 Laverne, OK 73848
 580-921-2029
 jolenag@health.ok.gov

OSDH Health Education
 Kathy Payne, Director
 1000 NE 10th St, room 506
 Oklahoma City, OK 73117
 405-271-6127
 KPayne@health.ok.gov

If you have an after-school program that is interested in learning more about CATCH Kids Club, a physical activity and nutrition program for children grades K-5, please contact the local health educator or Kathy Payne for information.

Primary Care Coverage Map


**Rate of Primary Care Physicians
per 100,000 Population, 2009 - 2010**



Note: The number of Primary Care Physicians for each county is represented under the county name.

Source: 2009-2010 Physician Survey, Office of Primary Care and Rural Health, Community Development Service, Oklahoma State Department of Health

OSDH Board of Health Map



Board of Health Members

- Alfred Baldwin, Jr.
- Barry L. Smith, J.D.
- Cris Hart-Wolfe (Treasurer)
- Jenny Alexopoulos, M.D. (President)
- Kenneth Miller, MD
- Michael D. Anderson, PhD (State at Large)
- R. Murali Krishna, M.D. (Vice-President)
- Richard G. Davis, DDS
- Ronald Woodson, MD

Created: 11.03.2010
Source: Oklahoma State Department of Health

Oklahoma Health Improvement Plan



For the complete OHIP, including a full list of partners, visit www.ok.gov/health and click the "Oklahoma Health Improvement Plan" link.



[STRATEGIC PLANNING]

FLAGSHIP GOALS

- Tobacco Use Prevention
- Obesity Reduction
- Children's Health

INFRASTRUCTURE GOALS

- Public Health Finance
- Workforce Development
- Access to Care
- Health Systems Effectiveness

SOCIETAL & POLICY INTEGRATION

- Policies and Legislation
- Social Determinants of Health & Health Equity

OKLAHOMA HEALTH IMPROVEMENT PLAN

Reference List

1. U.S. Census Bureau, Population Estimates, Accessed February 12, 2010 from www.census.gov.
2. U.S. Census Bureau, 2008 Census Estimates, Accessed February 12, 2010 from www.census.gov.
3. U.S. Census Bureau, Census 2000 Summary File 3, Accessed February 12, 2010 from www.census.gov
4. Oklahoma State Department of Health, Health Care Information, OK2SHARE, Death Statistics - Final, Accessed April 13, 2010, www.health.ok.gov/ok2share.
5. Oklahoma State Department of Health, Health Care Information, OK2SHARE, Inpatient Discharge Statistics, Accessed May 20, 2010, www.health.ok.gov/ok2share. Note: Data does not include federal or tribal facilities.
6. Sturm, R., 2002. The effects of obesity, smoking, and drinking on medical problems and costs. *Health Affairs*, 21, 245.
7. Oklahoma State Department of Health, Health Care Information, OK2SHARE, Youth Risk Behavior Survey (YRBS), www.health.ok.gov/ok2share.
8. Oklahoma State Department of Health, Health Care Information, OK2SHARE, Behavioral Risk Factor Surveillance Survey (BRFSS), www.health.ok.gov/ok2share.
9. Winning at Work—Diabetes Facts, American Diabetes Association, Accessed May 24, 2010 from www.diabetes.org
10. United Health Foundation, America's Health Rankings, www.americashealthrankings.org
11. Healthy Teen Network, Unique Needs of Children Born to Teen Parents, Accessed May 24, 2010 from www.healthteennetwork.org
12. Oklahoma State Department of Health, Health Care Information, OK2SHARE, Birth Statistics—Final, www.health.ok.gov/ok2share.
13. Than National Campaign to prevent Teen Pregnancy, By the Numbers: The Public Costs of Teen Childbearing in Oklahoma, November 2006.
14. Finch, B. K., 2003. Early origins of the gradient: The relationship between socioeconomic status and infant mortality in the United States. *Demography*, 40, 675-699.
15. United States Department of Health and Human Services (US DHHS), Centers of Disease Control and Prevention (CDC), National Center for Health Statistics (NCHS), Office of Analysis and Epidemiology (OAE), Division of Vital Statistics (DVS), Linked Birth / Infant Death Records 2003-2005 on CDC WONDER On-line Database. Accessed at <http://wonder.cdc.gov/lbd-current.html> on May 25, 2010 4:08:29 PM
16. National Safety Council, Estimating the Costs of Unintentional Injuries, Accessed May 25, 2010 from http://www.nsc.org/news_resources/injury_and_death_statistics/Pages/EstimatingtheCostsofUnintentionalInjuries.aspx.
17. March of Dimes, Low Birthweight Fact Sheet, Accessed May 25, 2010 from http://www.marchofdimes.com/professionals/14332_1153.asp#head2.
18. United States Department of Health and Human Services (US DHHS), Centers for Disease Control and Prevention (CDC), National Center for Health Statistics (NCHS), Division of Vital Statistics, Natality public-use data 2003-2006, on CDC WONDER Online Database, March 2009. Accessed at <http://wonder.cdc.gov/nativity-current.html> on May 25, 2010 5:09:49 PM
19. Oklahoma State Department of Health, 2005 State of the State's Health, Accessed May 25, 2010 from <http://www.ok.gov/health/pub/boh/state05/index.html>.
20. Oklahoma State Department of Health, Tobacco Use Prevention Service.
21. Centers for Disease Control and Prevention. Annual smoking-attributable mortality, years of potential life lost, and economic costs—United States, 1995–1999. *MMWR* 2002;51(14):300–303.



The Oklahoma Turning Point Initiative is public health improvement in action involving partnerships between the state and county departments of health, local communities, and policy-makers. The Oklahoma Turning Point engine is fueled by a community-based decision making process whereby local communities tap into the capacities, strengths, and vision of their citizens to create and promote positive, sustainable changes in the public health system, and the public’s health.

OKLAHOMA STATE DEPARTMENT OF HEALTH

Community and Family Health Services
 Community Development Service
 1000 NE 10th St, Room 508
 Oklahoma City, OK 73117
 Phone: 405-271-6127

Report compiled by:
 Miriam McGaugh, Ph.D.
 Epidemiologist
 and
 Shu-Chuan Lin, Ph.D.
 Epidemiologist

Working Together For Health

<http://www.ok.gov/health>

We are at a cross roads in our state and in Ellis County. Please come and be part of the solutions that will lead Oklahoma and Ellis County to becoming a healthy place to live, work and learn.

“If we are together nothing is impossible.
If we are divided all will fail.” - [Winston Churchill](#)

Turning Point Contact Information

If you are interested in learning more about Turning Point or becoming involved in local activities, please contact:

Lana Shaffer
 Harper County Health Department
 PO Box 290
 Laverne, OK 73848
 (580) 921-2029
 Email: LanaS@health.ok.gov
 Website: www.okturningpoint.org